

[Online First](#)

[Current Issue](#)

[Archive](#)


[Most Downloaded](#)

[Most Accessed](#)

[Indexing](#)

[Author Login](#)

[Reviewer Login](#)



International Journal of Medical Science and Public Health

ISSN (online) : 2277-338X

ISSN (Print) : 2320-4664


ISSUES : 12/year

PUBLISHER : Mrs. Dipika Charan

Micronutrients and growth of children: a literature review

Aminuddin Syam, Sukri Palutturi, Nurhaedar Djafar, Budu, Nurpudji Astuti, Abdul Razak Thaha

Int J Med Sci Public Health. 2016; 5(10): 1981-1986

[Online First](#)[Current Issue](#)[Archive](#)[Most Downloaded](#)[Most Accessed](#)[Indexing](#)[Author Login](#)[Reviewer Login](#)

International Journal of Medical Science and Public Health

ISSN (online) : 2277-338X

ISSN (Print) : 2320-4664

ISSUES : 12/year

PUBLISHER : Mrs. Dipika Charan

Micronutrients and growth of children: a literature review

Aminuddin Syam, Sukri Palutturi, Nurhaedar Djafar, Budu, Nurpudji Astuti, Abdul Razak Thaha

Int J Med Sci Public Health. 2016; 5(10): 1981-1986

Micronutrients and growth of children: A literature review

Aminuddin Syam^{1,2}, Sukri Palutturi³, Nurhaedar Djafar², Budu⁴, Nurpudji Astuti⁴, Abdul Razak Thaha⁵

¹Doctoral Program Student, Medical School Hasanuddin University, Makassar, Indonesia.

²Faculty of Public Health, Study Program of Nutritional Sciences, Hasanuddin University, Makassar, Indonesia.

³Faculty of Public Health, Department of Health Policy and Administration, Hasanuddin University, Makassar, Indonesia.

⁴Medical School of Hasanuddin University, Makassar, Indonesia.

⁵Faculty of Public Health, Nutritional Sciences, Hasanuddin University, Makassar, Indonesia.

Correspondence to: Aminuddin Syam, E-mail: amin.gzuh@gmail.com

Received June 10, 2016. Accepted August 12, 2016

Abstract

Optimal growth and development in children will be able to be achieved if the nutritional needs can be met. There are many teenagers who are malnourished because of inadequate intake. This study aims to examine the relationship between micronutrient especially folic acid and zinc with growth. The method used is a literature review of various sources of free accredited journals in pdf form eg PubMed, Proquest, Google Scholar and EBSCO. Other sources were such as books, reports national and international health. We collected literature published in the last 10 years to maintain the novelty of information. Some sources of literature were beyond the time limit if those are beneficial to this study. The results of the literature review indicate that there is a relationship between micronutrient and growth but this is still being debated. This study recommends for further follow-up especially on experimental studies in certain age groups.

KEY WORDS: Micronutrients, folic acid, zinc and growth

Introduction

Optimal growth and development in children will be able to achieve if the nutritional needs can be met. However, in reality, there are many teenagers who are malnourished due to inadequate intake and the presence of infectious diseases.^[1] Micronutrients, to the attention of many people especially public health experts and nutrition, are directly related to the growth and intelligence. Although, the data deficiency of zinc as an essential part of the micronutrient in Indonesia, for example, seems to be not yet fully available, but need to be aware of this possibility prevalence of zinc deficiency. A study conducted in the northeastern province of Thailand showed that about 70% of school children have low zinc levels and potentially affect their academic ability and growth.^[2,3] Furthermore, the lack of zinc can provide a wide range of impacts e.g. somatic growth of children and endocrine

systems as well as inhibition of sexual maturation, hypogonadism, and thyroid dysfunction.^[4]

Food and Agriculture Organization has set 4 strategies to solve the problem of micronutrient deficiencies, such as (1) the diversification of food, (2) food fortification, (3) supplementation with vitamins and minerals, as well as (4) the measurement of global public health and disease control.^[5] Food fortification is one of the best strategies because it has economic value with a higher level of compliance, so this way is more effective in reducing nutrient deficiency problems. In addition, fortification can also be used as a means of safe and effective way to supplement your diet with a low content of iron. Some countries have made efforts fortification in some products such as flour, oil, soy sauce, noodles, and rice.^[6,7]

In Indonesia, rice fortification can be a prospective strategy in addressing micronutrient deficiencies because rice is the main commodity. Almost all of Indonesia's population consumes rice. According to data Susenas in 2013, that the average consumption of rice per day of Indonesia's population reached 263.9 g/person, and not much difference between consumption in urban areas (253.3 g/person/day) and rural (274.4 g/person/day). Currently, the rice is only able to contribute calories but low in micronutrients such as folic acid, vitamin B1, niacin, iron, and zinc. Therefore, through the fortification of rice can provide micronutrient intake when consumed.^[8]

Based on the cost estimates by some experts that in addition to iron, which is known cost-effective in the fortification

Access this article online

Website: <http://www.ijmsph.com>

DOI: 10.5455/ijmsph.2017.10062016596

Quick Response Code:



- Sadeghi et al.^[25] To present data from the association of plasma zinc, copper and toxic elements of lead and cadmium levels with bone mineral density in Iranian women.
- No differences were found in the nutritional status, number of diseases, drugs, and functional activities between these groups. Plasma Zn, Cu, Pb, Cd levels were analyzed by a method of voltammetry. Mean \pm SD levels of copper and zinc was 1.168 ± 0.115 , 1.097 ± 0.091 $\mu\text{g/ml}$ in control group, 1.394 ± 0.133 , 1.266 ± 0.11 $\mu\text{g/ml}$ in total patient (TP) and 1.237 ± 0.182 , 1.127 ± 0.176 $\mu\text{g/ml}$ in mild patients ($-1 > \text{T-score} > -1.7$), 1.463 ± 0.174 , 1.327 ± 0.147 $\mu\text{g/ml}$ in severe patient group ($\text{T-score} < -1.7$); respectively. Mean \pm SD plasma level of lead and cadmium was 168.42 ± 9.61 ng/l , 2.91 ± 0.18 ng/ml in control group, 176.13 ± 8.64 ng/l , 2.97 ± 0.21 ng/ml in TP, 176.43 ± 13.2 ng/l , 2.99 ± 0.1 ng/ml in mild patients, 221.44 ± 20 ng/l and 3.80 ± 0.70 ng/ml in severe patient group, respectively. In this study plasma zinc, copper, lead, and cadmium concentrations were higher in the patients than in the control, though differences were not significant. However, differences were higher between the controls and patients with severe disease ($\text{T-score} < -1.7$). In addition adjusted T-score of femur with age and BMI showed negative significant correlation with plasma levels of zinc and lead in total participants ($p < 0.05$, $r = -0.201$, $p = 0.044$, $r = -0.201$).
- Shaikhkhalil et al.^[31] To evaluate the effect of enteral zinc supplementation on weight gain and linear growth in ELBW infants with CLD who received volume-restricted fortified human milk.
- Zinc supplementation started at postmenstrual age 33 ± 2 weeks. Most infants received fortified human milk. Weight gain increased from 10.9 before supplementation to 19.9 $\text{g kg}^{-1} \text{ day}^{-1}$ after supplementation ($p < 0.0001$). Linear growth increased from 0.7 to 1.1 cm/week ($p = 0.001$). Zinc supplementation improved growth in ELBW infants with CLD receiving human milk. Further investigation is warranted to re-evaluate zinc requirements, markers, and balance.

Acknowledgement

This study is part of a study of literature that has been adapted from my Doctor Program at the Medical Faculty of Hasanuddin University, Indonesia. Therefore, our gratitude goes to the promoter and co-promoter as well as to the Indonesian government who have helped in completing this study.

References

- UNICEF. Conceptual Framework for Nutritional Status 1998.
- Black MM. Zinc deficiency and child development. *Am J Clin Nutr* 1998;68(2):464S-9S.
- Thurlow R, Winichagoon P, Pongcharoen T, Gowachirapant S, Boonpradern A, Manger M, et al. Risk of zinc, iodine and other micronutrient deficiencies among school children in North East Thailand. *Euro J Clin Nutr* 2006;60(5):623-32.
- Laitinen R. Zinc, copper, and sexual maturation in 9-18-year-old girls and boys. *Biological Trace Ele Res* 1990;25(1):71-8.
- Hussain A. Preventing and controlling micronutrient malnutrition through food-based actions in South Asian countries. Food, Nutrition and Agriculture (FAO) Alimentation, Nutrition et Agriculture (FAO) Alimentacion, Nutricion y Agricultura (FAO) 1998.
- Nesamvuni AE, Vorster HH, Margetts BM, Kruger A. Fortification of maize meal improved the nutritional status of 1-3-year-old African children. *Pub Health Nutr* 2005;8(05):461-7.
- Theary C, Panagides D, Laillou A, Vonthanak S, Kanarath C, Chhorvann C, et al. Fish sauce, soy sauce, and vegetable oil fortification in Cambodia: where do we stand to date? *Food Nutr Bull* 2013;34(2 suppl1):S62-S71.
- Radhika MS, Nair KM, Kumar RH, Rao MV, Ravinder P, Reddy CG, et al. Micronized ferric pyrophosphate supplied through extruded rice kernels improves body iron stores in children: a double-blind, randomized, placebo-controlled midday meal feeding trial in Indian schoolchildren. *Am J Clin Nutr* 2011;94(5):1202-10.
- Romano PS, Waitzman NJ, Scheffler RM, Pi RD. Folic acid fortification of grain: an economic analysis. *Am J Pub Health* 1995;85(5):667-76.
- IZINCG IZINCG. Zinc fortification 2007.
- Soliman AT, Sanctis VD, Kalra2 S. Anemia and Growth. *Ind J Endocrinol Metabol* 2014;18:S1-S5.
- World Health Organization. Iron deficiency anaemia: assessment, prevention and control: a guide for programme managers, 2001.
- World Health Organization. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity, Available 2011.
- Sen A, Kanani SJ. Impact of iron-folic acid supplementation on cognitive abilities of school girls in Vadodara. *Ind Pediat* 2009;46(2):137.
- Katelhut A, Schultink W, Angeles I, Gross R, Pietrzik K. The effects of weekly iron supplementation with folic acid, vitamin A, vitamin C on iron status of Indonesian adolescents. *Asia Pac J Clin Nutr* 1996;5(3):181-5.
- Sen A, Kanani S. Intermittent iron folate supplementation: impact on hematinic status and growth of school girls. *ISRN hematology* 2012;2012.

- Sadeghi et al.^[25] To present data from the association of plasma zinc, copper and toxic elements of lead and cadmium levels with bone mineral density in Iranian women.
- No differences were found in the nutritional status, number of diseases, drugs, and functional activities between these groups. Plasma Zn, Cu, Pb, Cd levels were analyzed by a method of voltammetry. Mean \pm SD levels of copper and zinc was 1.168 ± 0.115 , 1.097 ± 0.091 $\mu\text{g/ml}$ in control group, 1.394 ± 0.133 , 1.266 ± 0.11 $\mu\text{g/ml}$ in total patient (TP) and 1.237 ± 0.182 , 1.127 ± 0.176 $\mu\text{g/ml}$ in mild patients ($-1 > \text{T-score} > -1.7$), 1.463 ± 0.174 , 1.327 ± 0.147 $\mu\text{g/ml}$ in severe patient group ($\text{T-score} < -1.7$); respectively. Mean \pm SD plasma level of lead and cadmium was 168.42 ± 9.61 ng/l , 2.91 ± 0.18 ng/ml in control group, 176.13 ± 8.64 ng/l , 2.97 ± 0.21 ng/ml in TP, 176.43 ± 13.2 ng/l , 2.99 ± 0.1 ng/ml in mild patients, 221.44 ± 20 ng/l and 3.80 ± 0.70 ng/ml in severe patient group, respectively. In this study plasma zinc, copper, lead, and cadmium concentrations were higher in the patients than in the control, though differences were not significant. However, differences were higher between the controls and patients with severe disease ($\text{T-score} < -1.7$). In addition adjusted T-score of femur with age and BMI showed negative significant correlation with plasma levels of zinc and lead in total participants ($p < 0.05$, $r = -0.201$, $p = 0.044$, $r = -0.201$).
- Shaikhkhalil et al.^[31] To evaluate the effect of enteral zinc supplementation on weight gain and linear growth in ELBW infants with CLD who received volume-restricted fortified human milk.
- Zinc supplementation started at postmenstrual age 33 ± 2 weeks. Most infants received fortified human milk. Weight gain increased from 10.9 before supplementation to 19.9 $\text{g kg}^{-1} \text{ day}^{-1}$ after supplementation ($p < 0.0001$). Linear growth increased from 0.7 to 1.1 cm/week ($p = 0.001$). Zinc supplementation improved growth in ELBW infants with CLD receiving human milk. Further investigation is warranted to re-evaluate zinc requirements, markers, and balance.

Acknowledgement

This study is part of a study of literature that has been adapted from my Doctor Program at the Medical Faculty of Hasanuddin University, Indonesia. Therefore, our gratitude goes to the promoter and co-promoter as well as to the Indonesian government who have helped in completing this study.

References

- UNICEF. Conceptual Framework for Nutritional Status 1998.
- Black MM. Zinc deficiency and child development. *Am J Clin Nutr* 1998;68(2):464S-9S.
- Thurlow R, Winichagoon P, Pongcharoen T, Gowachirapant S, Boonpradern A, Manger M, et al. Risk of zinc, iodine and other micronutrient deficiencies among school children in North East Thailand. *Euro J Clin Nutr* 2006;60(5):623-32.
- Laitinen R. Zinc, copper, and sexual maturation in 9-18-year-old girls and boys. *Biological Trace Ele Res* 1990;25(1):71-8.
- Hussain A. Preventing and controlling micronutrient malnutrition through food-based actions in South Asian countries. Food, Nutrition and Agriculture (FAO) Alimentation, Nutrition et Agriculture (FAO) Alimentacion, Nutricion y Agricultura (FAO) 1998.
- Nesamvuni AE, Vorster HH, Margetts BM, Kruger A. Fortification of maize meal improved the nutritional status of 1-3-year-old African children. *Pub Health Nutr* 2005;8(05):461-7.
- Theary C, Panagides D, Laillou A, Vonthanak S, Kanarath C, Chhorvann C, et al. Fish sauce, soy sauce, and vegetable oil fortification in Cambodia: where do we stand to date? *Food Nutr Bull* 2013;34(2 suppl1):S62-S71.
- Radhika MS, Nair KM, Kumar RH, Rao MV, Ravinder P, Reddy CG, et al. Micronized ferric pyrophosphate supplied through extruded rice kernels improves body iron stores in children: a double-blind, randomized, placebo-controlled midday meal feeding trial in Indian schoolchildren. *Am J Clin Nutr* 2011;94(5):1202-10.
- Romano PS, Waitzman NJ, Scheffler RM, Pi RD. Folic acid fortification of grain: an economic analysis. *Am J Pub Health* 1995;85(5):667-76.
- IZINCG IZINCG. Zinc fortification 2007.
- Soliman AT, Sanctis VD, Kalra2 S. Anemia and Growth. *Ind J Endocrinol Metabol* 2014;18:S1-S5.
- World Health Organization. Iron deficiency anaemia: assessment, prevention and control: a guide for programme managers, 2001.
- World Health Organization. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity, Available 2011.
- Sen A, Kanani SJ. Impact of iron-folic acid supplementation on cognitive abilities of school girls in Vadodara. *Ind Pediat* 2009;46(2):137.
- Katelhut A, Schultink W, Angeles I, Gross R, Pietrzik K. The effects of weekly iron supplementation with folic acid, vitamin A, vitamin C on iron status of Indonesian adolescents. *Asia Pac J Clin Nutr* 1996;5(3):181-5.
- Sen A, Kanani S. Intermittent iron folate supplementation: impact on hematinic status and growth of school girls. *ISRN hematology* 2012;2012.